

Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) EP 1 227 088 B1

(12) EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
17.09.2003 Bulletin 2003/38

(51) Int Cl.7: C07D 307/87, A61K 31/343,
A61P 25/24

(21) Application number: 02009350.6

(22) Date of filing: 28.02.2001

(54) Crystalline base of citalopram and hydrochloride or hydrobromide salt thereof
Kristalline Base von Citalopram, und Hydrochlorid- oder Hydrobromidsalz davon
Base cristalline de citalopram et son sel hydrochlorique ou hydrobromique

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: 13.03.2000 DK 200000402
13.04.2000 WOPCT/DK00/00183

(43) Date of publication of application:
31.07.2002 Bulletin 2002/31

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
01909568.6 / 1 169 314

(73) Proprietor: H.Lundbeck A/S
2500 Valby-Copenhagen (DK)

(72) Inventors:
• Petersen, Hans
2720 Vanløse (DK)
• Bögesø, Klaus Peter
2970 Hørsholm (DK)
• Holm, Per
2720 Vanløse (DK)

(74) Representative: HOFFMANN - EITLE
Patent- und Rechtsanwälte
Arabellastrasse 4
81925 München (DE)

(56) References cited:
EP-A- 0 171 943 EP-A- 0 347 066
WO-A-98/19511 WO-A-98/19512
WO-A-98/19513 DE-A- 2 657 013
DE-U- 20 007 303

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).


Printed by Jouve, 75001 PARIS (FR)

EP 1 227 088 B1

BEST AVAILABLE COPY

Claims

1. Crystalline base of citalopram, characterised in that it has a purity of more than 99.8 %w/w, preferably more than 99.9 % w/w.
2. Crystalline base of citalopram, prepared by a process characterised in that the base of citalopram is set free and precipitated in crystalline form and optionally re-crystallised one or more times.
3. The crystalline base of citalopram according to claim 2 characterised in that the base of citalopram is set free from a crude salt or a crude mixture of citalopram.
4. Crystalline base of citalopram, prepared by a process which is characterised in that one or more impurities of the formula




wherein Z is halogen, $-O-SO_2-(CF_2)_n-CF_3$, where n is 0-8, $-CHO$, $-NHR^1$, $-COOR^2$, $-CONR^2R^3$ wherein R^2 and R^3 are selected from hydrogen, alkyl, optionally substituted aryl or aralkyl and R^1 is hydrogen or alkylcarbonyl, are removed from a crude mixture of citalopram or from a crude salt of citalopram, by precipitating citalopram base in crystalline form and optionally re-crystallising said base one or more times.

5. The crystalline base of citalopram according to claim 4 wherein the crude mixture of citalopram containing the compound of formula II as an impurity, is prepared by subjecting a compound of formula II to a cyanide exchange reaction with a cyanide source.
6. The crystalline base of citalopram according to claim 4 wherein Z is halogen, in particular bromide or chloride.
7. The crystalline base of citalopram according to any one of claims 4 to 6 wherein the crude mixture of citalopram is subjected to initial purification before the base of citalopram is precipitated in crystalline form.
8. The crystalline base of citalopram according to any one of claims 4 to 6 wherein the crude mixture of citalopram is subjected to initial purification before a crude salt is formed from said crude mixture.
9. The crystalline base of citalopram according to any one of claims 4 to 7 wherein the base of citalopram is set free from a crude salt or a crude mixture of citalopram by treatment with a base and optionally subjected to further purification before the base of citalopram is precipitated in crystalline form.
10. The crystalline base of citalopram according to any one of claims 2 to 4, characterised in that the crude salt is a hydrobromide, hydrochloride, sulphate, oxalate, phosphate or nitrate salt, preferably the sulphate, hydrobromide, or hydrochloride salt.
11. The crystalline base of citalopram according to any one of claims 2 to 10, characterised in that it has a purity of more than 99.8 %w/w, preferably more than 99.9 %w/w.
12. A pharmaceutical composition containing the crystalline base of citalopram according to any one of claims 1 to 11.
13. The pharmaceutical composition according to claim 12 which is a tablet prepared by

= 12.13.2003

= C



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 227 088 B1**

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention
of the grant of the patent:
17.09.2003 Bulletin 2003/38

(51) Int Cl.7: **C07D 307/87, A61K 31/343,
A61P 25/24**

(21) Application number: **02009350.6**

(22) Date of filing: **28.02.2001**

(54) **Crystalline base of citalopram and hydrochloride or hydrobromide salt thereof**

Kristalline Base von Citalopram, und Hydrochlorid- oder Hydrobromidsalz davon

Base crystalline de citalopram et son sel hydrochlorique ou hydrobromique

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **13.03.2000 DK 200000402
13.04.2000 WOPCT/DK00/00183**

(43) Date of publication of application:
31.07.2002 Bulletin 2002/31

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
01909568.6 / 1 169 314

(73) Proprietor: **H.Lundbeck A/S
2500 Valby-Copenhagen (DK)**

(72) Inventors:
• **Petersen, Hans
2720 Vanløse (DK)**
• **Bøgesø, Klaus Peter
2970 Hørsholm (DK)**
• **Holm, Per
2720 Vanløse (DK)**

(74) Representative: **HOFFMANN - EITLE
Patent- und Rechtsanwälte
Arabellastrasse 4
81925 München (DE)**

(56) References cited:
**EP-A- 0 171 943 EP-A- 0 347 066
WO-A-98/19511 WO-A-98/19512
WO-A-98/19513 DE-A- 2 657 013
DE-U- 20 007 303**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).


EP 1 227 088 B1

Description

[0001] The present invention relates to the crystalline base of the well known antidepressant drug citalopram, 1-[3-(dimethylamino)propyl]-1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofurancarbonitrile and formulations of said base.

Background of the Invention

[0002] Citalopram is a well-known antidepressant drug that has now been on the market for some years and has the following structure:



[0003] It is a selective, centrally acting serotonin (5-hydroxytryptamine; 5-HT) reuptake inhibitor, accordingly having antidepressant activities. The antidepressant activity of the compound has been reported in several publications, eg. J. Hyttel, *Prog. Neuro-Psychopharmacol. & Biol. Psychiat.*, 1982, 6, 277-295 and A. Gravem, *Acta Psychiatr. Scand.*, 1987, 75, 478-486. The compound has further been disclosed to show effects in the treatment of dementia and cerebrovascular disorders, EP-A-474580.

[0004] Citalopram was first disclosed in DE 2,657,013, corresponding to US 4,136,193. This patent publication describes the preparation of citalopram by one method and outlines a further method, which may be used for preparing citalopram. The citalopram prepared was isolated as the oxalate, the hydrobromide and the hydrochloride salt, respectively. Furthermore, the citalopram base was obtained as an oil (B.P. 175 C/0.03 mmHg). Citalopram is marketed as the hydrobromide and the hydrochloride, respectively.

[0005] A number of processes for the preparation of citalopram have been disclosed. In many of these, the last step of the process is a conversion of a group different from cyano in the 5 position of the direct analogue of citalopram to a 5-cyano group. So citalopram has been prepared by:

Exchange of 5-halogen, or 5-CF₃-(CF₂)_n-SO₂-O- with cyano (DE 2,657,013 and co-pending WO 0011926 and WO 0013648)

Conversion of a 5-amido or 5-ester group to a 5-cyano group (WO 9819513)

Conversion of a 5-amino group to a 5-cyano group (WO 9819512)

Conversion of a 5-formyl group to a 5-cyano group (WO 9900548)

Conversion of a 5-oxazoliny or 5-thiazoliny group to a 5-cyano group (WO 0023431)

[0006] Other processes for the preparation of citalopram comprise exchange of the 5-bromo group of 1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofuranbromide with 5-cyano followed by alkylation with a 3-(N,N-dimethylamino)propyl-halogenide (DE 2,657,013 and WO 9819511).

[0007] Many of the processes mentioned above have the disadvantage that it is difficult to separate the intermediates formed during the process (the intermediates mentioned above or earlier intermediates) from the end product and, accordingly, extensive purification procedures involving loss of citalopram are required in order to obtain the necessary quality of the end product.


[0008] It has now been found that the base of citalopram may be obtained as a very nice and pure crystalline product, which may easily be handled and conveniently be formulated into tablets and other pharmaceutical forms. Furthermore, it has surprisingly been found that a very good and efficient purification of citalopram may be obtained during manufacture of citalopram (e.g. of the hydrobromide or the hydrochloride salt) by crystallising the base, and thereafter optionally forming a salt from the base.

[0009] This purification process is particularly useful for removing intermediates which are structurally closely related to citalopram, in particular compounds which only differ from citalopram by the substituent situated in position 5 on the


isobenzofurane ring, and intermediates which have physical/chemical properties which are close to those of citalopram, e.g. the 1-[3-(dimethylamino)propyl]-1-(4-fluorophenyl)-1,3-dihydro-isobenzofuranes having halogen (in particular bromide and chloride), an amide or an ester in position 5 of the isobenzofurane ring, or 1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofuranbromide, or -chloride.

Summary of the invention

[0010] The present invention provides the crystalline base of the compound



[0011] More particularly, the present invention relates to a process for the manufacture of citalopram base characterised in that one or more impurities of the formula



wherein Z is halogen, $-O-SO_2-(CF_2)_n-CF_3$, where n is 0-8, $-CHO$, $-NHR^1$, $-COOR^2$, $-CONR^2R^3$ wherein R^2 and R^3 are selected from hydrogen, alkyl, optionally substituted aryl or aralkyl and R^1 is hydrogen or alkylcarbonyl, are removed from a crude mixture of citalopram or from a crude salt of citalopram, by precipitating citalopram base in crystalline form, and optionally re-crystallising said base one or more times.

[0012] The crude mixture of citalopram containing the compound of formula II as an impurity may be prepared by subjecting a compound of formula II to a cyanide exchange reaction with a cyanide source, or by subjecting 1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofuranhalogenide, in particular the bromide, to a cyanide exchange reaction followed by alkylation with a 3-(N,N-dimethylamino)propyl-halogenide.

[0013] In a particular embodiment of the invention, Z is halogen, in particular bromide or chloride.

[0014] The crude salt may be any convenient salt, such as the hydrobromide, hydrochloride, sulphate, oxalate, phosphate, nitrate or any other convenient salt. Other salts are salts of organic acids.

[0015] In a preferred embodiment of the invention, the crude salt is the sulphate, the hydrobromide or the hydrochloride salt.

[0016] In yet another aspect, a pharmaceutical formulation of the free base of citalopram is provided. Preferably the formulation is for oral administration.

[0017] The formulations according to the invention may be prepared by direct compression of citalopram in admixture with conventional adjuvants or diluents. Alternatively, a wet granulate or a melt granulate of citalopram, optionally in admixture with conventional adjuvants or diluents may be used for compression of tablets.

[0018] In particular, the pharmaceutical composition of the invention contains the racemic mixture of citalopram base.

[0019] The crystalline base of citalopram is preferably more than 99.8% w/w pure, most preferably more than 99.9%

w/w pure (peak area). The melting point is preferably a range within 90 - 93 °C, most preferably 91 - 92 °C (DSC; onset, open capsule) or it is between 92 and 94°C, preferably 92.5 and 93.5 °C (DSC; onset, closed capsule). The crystalline base of citalopram is preferably in racemic form.

[0020] The terms "crude salt" and "crude mixture" refer to the fact that the salt and the mixture, respectively, comprise impurities, in particular impurities of formula II, which must be removed or which it is desired to remove.

[0021] The crude salt may be a salt separated directly from the reaction mixture, or the crude reaction mixture may have been subjected to some initial purification, e.g. one re-crystallisation, and/or treatment with activated carbon or silica gel, and the salt formed subsequently by treatment with an acid using methods known in the art. The salt may be isolated by precipitation or it may exist in a solvent, e.g. in the mixture resulting directly from the synthesis of the salt.

[0022] Similarly, the crude mixture comprising citalopram may be obtained directly from the synthesis of the compound according to any of the above mentioned processes or it may have been subjected to some initial or simultaneous purification, e.g. one re-crystallisation, treatment with activated carbon or silica gel.

[0023] The base of citalopram may be set free from the crude salt by dissolving the crude salt in a mixture of water and an organic solvent and then adding a base. The organic solvent may be toluene, ethyl acetate or any other suitable solvent and the base may be any convenient base, preferably NaOH or NH₃. Likewise, the base of citalopram may, if necessary, be set free from a crude mixture containing citalopram by treatment with a base.

[0024] Crude mixtures containing citalopram base may be subjected to further purification and extraction, before the base is precipitated in crystalline form. The base of citalopram may be isolated by separation of the organic phase, evaporation of the solvent in order to obtain the base most probably as an oil and then crystallisation of the base from an aprotic solvent, such as an alkane, including n-heptane, hexane and isooctane, and high and low boiling petroleum ethers and substituted aromates, incl toluene and xylenes. Crystalline citalopram base may be re-crystallised from the same solvents.

[0025] The cyanide exchange reactions mentioned above may be carried out as described in the patent applications mentioned above.

[0026] In particular, when Z is halogen, or CF₃-(CF₂)_n-SO₂-O- wherein n is an integer in the range 0-8, incl., the conversion to a cyano group may be carried out by reaction with a cyanide source, for example KCN, NaCN, CuCN, Zn(CN)₂ or (R⁴)₄NCN where R⁴ indicates four groups which may be the same or different and are selected from hydrogen and straight chain or branched alkyl, in the presence of a palladium catalyst and a catalytic amount of Cu⁺ or Zn²⁺, or with Zn(CN)₂ in the presence of a palladium catalyst.


[0027] The cyanide source is used in a stoichiometric amount or in excess, preferably 1-2 equivalents are used per equivalent starting material. (R⁴)₄N⁺ may conveniently be (Bu)₄N⁺. The cyanide compound is preferably NaCN or KCN or Zn(CN)₂.

[0028] The palladium catalyst may be any suitable Pd(0) or Pd(II) containing catalyst, such as Pd(PPh₃)₄, Pd₂(dba)₃, Pd(PPh)₂Cl₂, etc. The Pd catalyst is conveniently used in an amount of 1-10, preferably 2-6, most preferably about 4-5 mol%.

[0029] Catalytic amounts of Cu⁺ and Zn²⁺, respectively, means substoichiometric amounts such as 0.1- 5, preferably 1 - 3 %. Conveniently, about ½ eq. is used per eq. Pd. Any convenient source of Cu⁺ and Zn²⁺ may be used. Cu⁺ is preferably used in the form of CuI and Zn²⁺ is conveniently used as the Zn(CN)₂ salt.

[0030] When Z is Br or I, the conversion to a cyano group may also be carried out by reaction with Cu(CN) without catalyst. In a preferred embodiment, the reaction is performed at elevated temperature.

[0031] In another aspect the reaction is performed in an ionic liquid of the general formula (R⁵)₄N⁺, X⁻, wherein R⁵ are alkyl-groups or two of the R⁵ groups together form a ring and X⁻ is the counterion. In one embodiment, (R⁵)₄N⁺X⁻ represents



[0032] In another particular aspect, the reaction is conducted with apolar solvents such as benzene, xylene or mesitylene and under the influence of microwaves by using *i.e.* Synthewave 1000™ by Prolabo. In a particular aspect, the reaction is performed without added solvent.

[0033] The temperature ranges are dependent upon the reaction type. If no catalyst is present, preferred temperatures are in the range of 100-200°C. However, when the reaction is conducted under the influence of microwaves, the temperature in the reaction mixture may raise to above 300 °C. More preferred temperature ranges are between

120-170°C. The most preferred range is 130-150°C. If catalyst is present, the preferred temperature range is between 0 and 100°C. More preferred are temperature ranges of 40-90°C. Most preferred temperature ranges are between 60-90°C.

[0034] Other reaction conditions, solvents, etc. are conventional conditions for such reactions and may easily be determined by a person skilled in the art.

[0035] When Z is Cl or Br, the conversion to a cyano group may also be carried out by reaction with a cyanide source, for example KCN, NaCN, CuCN, Zn(CN)₂ or (R⁴)₄NCN where (R⁴)₄ indicates four groups which may be the same or different and are selected from hydrogen and straight chain or branched alkyl, in the presence of a nickel catalyst.

[0036] The nickel catalyst may be any suitable Ni(0) or Ni(II) containing complex which acts as a catalyst, such as Ni(PPh₃)₃, (σ-aryl)-Ni(PPh₃)₂Cl, etc. The nickel catalysts and their preparation are described in WO 96/11906, EP-A-613720 or EP-A-384392.

[0037] In one embodiment, the reaction is carried out in the presence of a catalytic amount of Cu⁺ or Zn²⁺.

[0038] In a particularly preferred embodiment, a Nickel(0) complex is prepared *in situ* before the cyanation reaction by reduction of a Nickel(II) precursor such as NiCl₂ or NiBr₂ by a metal, such as zinc, magnesium or mangan in the presence of excess of complex ligands, preferably triphenylphosphine.

[0039] The Ni-catalyst is conveniently used in an amount of 0.5-10, preferably 2-6, most preferably about 4-5 mol%. Catalytic amounts of Cu⁺ and Zn²⁺, respectively, mean substoichiometric amounts such as 0.1-5, preferably 1-3%. Any convenient source of Cu⁺ and Zn²⁺ may be used. Cu⁺ is preferably used in the form of CuI and Zn²⁺ is conveniently used as the Zn(CN)₂ salt or formed *in situ* by reduction of a Nickel(II) compounds using zinc.

[0040] The Ni catalysts are *i.e.* Ni(0), Pd(0) or Pd(II) catalysts as described by Sakakibara et. al. in Bull. Chem. Soc. Jpn., 61, 1985-1990, (1988). Preferred catalysts are Ni(PPh₃)₃ or Pd(PPh₃)₄, or Pd(PPh)₂Cl₂.

[0041] The reactions may be performed in any convenient solvent as described in Sakakibara et. al. in Bull. Chem. Soc. Jpn., 61, 1985-1990, (1988). Preferred solvents are acetonitrile, ethylacetate, THF, DMF or NMP.

[0042] When Z is CHO, the conversion to a cyano group may be carried out by conversion of the formyl group to an oxime or similar group by reaction with a reagent R⁶-V-NH₂ wherein R⁶ is hydrogen, optionally substituted alkyl, aryl or heteroaryl and V is O, N or S, followed by dehydration with a common dehydrating agent, for example thionylchloride, acetic anhydride/pyridine, pyridine/HCl or phosphor pentachloride. Preferred reagents R⁶-V-NH₂ are hydroxylamine and compounds wherein R⁶ is alkyl or aryl and V is N or O.

[0043] When Z is -COOH, the conversion to a cyano group may be carried out via the corresponding acid chloride, ester or amide.

[0044] The acid chloride is conveniently obtained by treatment of the acid with POCl₃, PCl₅ or SOCl₂ neat or in a suitable solvent, such as toluene or toluene comprising a catalytic amount of N,N-dimethylformamide. The ester is obtained by treatment of the acid with an alcohol, in the presence of an acid, preferably a mineral acid or a Lewis acid, such as HCl, H₂SO₄, POCl₃, PCl₅ or SOCl₂. Alternatively, the ester may be obtained from the acid chloride by reaction with an alcohol. The ester or the acid chloride is then converted to an amide by amidation with ammonia or an alkylamine, preferably *t*-butyl amine.

[0045] The conversion to amide may also be obtained by reaction of the ester with ammonia or an alkylamine under pressure and heating.

[0046] The amide group is then converted to a cyano group by dehydration. The dehydrating agent may be any suitable dehydrating agent, and the optimal agent may easily be determined by a person skilled in the art. Examples of suitable dehydrating agents are SOCl₂, POCl₃ and PCl₅, preferably SOCl₂.

[0047] In a particularly preferred embodiment, the carboxylic acid is reacted with an alcohol, preferably ethanol, in the presence of POCl₃, in order to obtain the corresponding ester, which is then reacted with ammonia thereby giving the corresponding amide, which in turn is reacted with SOCl₂ in toluene comprising a catalytic amount of N,N-dimethylformamide.

[0048] Alternatively, a compound where Z is -COOH may be reacted with chlorosulfonyl isocyanate in order to form the nitrile, or treated with a dehydrating agent and a sulfonamide.

[0049] When Z is -NHR¹, where R¹ is hydrogen, the conversion into cyano is preferably performed by diazotation and followed by reaction with CN⁻. Most preferably NaNO₂ and CuCN and/or NaCN are used. When R¹ is alkylcarbonyl, it is initially subjected to hydrolysis thereby obtaining the corresponding compound wherein R¹ is H which is then converted as described above. The hydrolysis may be performed either in acidic or basic environment.

[0050] The compounds of formula (II) may be prepared as described in DE 2,657,013, WO 0011926 and WO 0013648, WO 9819513, WO 9819512 and WO 9900548.

[0051] Throughout this specification with claims halogen means chloro, bromo or iodo.

[0052] The term alkyl refers to a branched or unbranched alkyl group, such as methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2-methyl-2-propyl, and 2-methyl-1-propyl.

[0053] The term aryl refers to a carbocyclic aromatic group, in particular phenyl. Arylalkyl refers to an arylalkyl group wherein aryl and alkyl is as defined above. The aryl and arylalkyl groups may optionally be substituted, e.g. with alkyl

groups, forming for example tolyl.

[0054] The pharmaceutical compositions of the invention may be administered in any suitable way and in any suitable form, for example orally in the form of tablets, capsules, powders or syrups, or parenterally in the form of usual sterile solutions for injection. Preferably the pharmaceutical compositions of the invention are administered orally.

[0055] The pharmaceutical formulations of the invention may be prepared by conventional methods in the art. For example, tablets may be prepared by mixing the active ingredient with ordinary adjuvants and/or diluents and subsequently compressing the mixture in a conventional tableting machine. Examples of adjuvants or diluents comprise: Corn starch, potato starch, talcum, magnesium stearate, gelatine, lactose, gums, and the like. Any other adjuvant or additive, colourings, aroma, preservatives etc. may be used provided that they are compatible with the active ingredients.

[0056] In particular, the formulations according to the invention may be prepared by direct compression of citalopram in admixture with conventional adjuvants or diluents. Alternatively, a wet granulate or a melt granulate of citalopram, optionally in admixture with conventional adjuvants or diluents may be used for compression of tablets.

[0057] Solutions for injections may be prepared by solving the active ingredient and possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution to the desired volume, sterilisation of the solution and filling in suitable ampoules or vials. Any suitable additive conventionally used in the art may be added, such as tonicity agents, preservatives, antioxidants, etc.

[0058] According to the present invention, the base of citalopram has been found to be crystalline with stable and nice white crystals and it has been found that the base may easily be crystallised in a very pure form. So for example more than 99.8% w/w pure citalopram base was obtained by crystallisation from up to 95% pure hydrobromide without further purification. Accordingly, the yield may be improved substantially during the manufacture of citalopram.

[0059] Finally, it has been found that the crystalline citalopram base may be formulated into very good and stable solid formulations with good release properties.

[0060] The invention is further illustrated by the following examples.

Example 1

Crystallisation of R,S-Citalopram as the free base.

[0061] 1-(3-Dimethylaminopropyl)-1-(4'-fluorophenyl)-1,3-dihydrobenzofuran-5-carbonitrile.

[0062] 1-(3-Dimethylaminopropyl)-1-(4'-fluorophenyl)-1,3-dihydrobenzofuran-5-carbonitrile hydrobromide (101 grams, 0.25 mole) prepared from 1-(3-Dimethylaminopropyl)-1-(4'-fluorophenyl)-1,3-dihydro-5-isobenzofuranbromide, is suspended in water (500 ml) and toluene (500 ml). NaOH (60 ml, 5 N (aq)) is added and the mixture (pH>10) is stirred for 15 min. before the phases are separated. The organic phase is washed with water (2x100 ml) and filtered through a pad of filter help. The volatiles are removed *in vacuo* and the title compound is obtained as an oil. n-Heptane (400 ml) is added and the mixture is heated to 70 °C. On cooling, crystals form. The white crystals of the title compound are filtered off and dried at ambient temperature over night *in vacuo*. Yield: 75.4 grams (93%). DSC(onset, open capsule): 91.3-91.8 °C DSC (onset, closed capsule): 92.8 °C. Purity: (> 99.8 % (peak area)). Anal. calcd. for C₂₀H₂₁N₂F₁O₁; C, 74.04; H, 6.54; N, 8.64. Found C, 74.01; H, 6.49; N, 8.59. ¹H-NMR (DMSO-d₆, 500 MHz): 1.21 (1H, m), 1.29 (1H, m), 2.02 (6H, s), 2.09-2.23 (4 H, m), 5.15 (1H, d J=12.5 Hz), 5.22 (1H, d J=12.5 Hz), 7.16 (2H, t J=8.5 Hz), 7.60 (2H, dt J=8.5 Hz J=1.2 Hz), 7.76 (1H, d J= 8.5 Hz), 7.79 (1H, d J=8.5 Hz), 7.80 (1H, s). ¹³C-NMR (DMSO-d₆, 125 MHz): 21.8, 38.3, 45.0, 58.8, 71.0, 90.7, 110.5, 115.1 (d J=22 Hz), 118.8, 123.1, 125.1, 127.0 (d J=8 Hz), 132.0, 140.0 (d J=3 Hz), 140.5, 149.5, 161.3 (d J=245Hz).

Example 2

[0063]

a) A crude mixture of Citalopram and sulphuric acid is made basic by adding NaOH and the citalopram base is extracted with toluene. The toluene is evaporated and the citalopram base obtained is dissolved in n-heptane at elevated temperature. The very pure free base of citalopram is precipitated by cooling.

b) A crude mixture of Citalopram and sulphuric acid is made basic by adding NaOH and the citalopram base is extracted with toluene. The toluene is evaporated and the citalopram base obtained is dissolved in methanol. The mixture is treated with activated carbon and filtrated and the solvent is evaporated. The purified free base is dissolved in n-heptane at elevated temperature. Then the very pure free base of citalopram is precipitated by cooling.

c) A crude mixture of Citalopram and sulphuric acid is made basic by adding NaOH and the citalopram base is

EP 1 227 088 B1

extracted with toluene. The toluene phase is treated with silicagel, the toluene is evaporated and the citalopram base obtained is dissolved in n-heptane at elevated temperature. The very pure free base of citalopram is precipitated by cooling.

d) A crude mixture of Citalopram and sulphuric acid is made basic by adding NaOH and the citalopram base is extracted with toluene. The toluene phase is treated with silicagel, the toluene is evaporated and the citalopram base obtained is dissolved in methanol. The mixture is treated with activated carbon and filtrated and the solvent is evaporated. The purified free base is dissolved in n-heptane at elevated temperature. Then the extremely pure free base of citalopram is precipitated by cooling.

Example 3

Wet granulation and preparation of tablets

[0064] The batch size was 200 g and the granulation was performed in a small-scale laboratory high shear mixer (Micromixer).

[0065] Citalopram base was sieved through a sieve aperture of 0.3 mm. The ingredients of the intragranular phase (1 - 4 in Table 1) were mixed at 600 rpm. 25 ml of purified water (5) was added in 30 sec and the granulation terminated after a total processing time of 3 min. The granulate was wet sieved through a 0.7 mm sieve aperture and dried at 40 °C in 30 minutes to equilibrium relative humidity of 32 %. The dried granulate was finally sieved through a 0.7 mm sieve aperture.

[0066] The dried granulate was mixed for 3 minutes with the extragranular phase (6 - 7) in a Turbula mixer and finally mixed with the lubricant (8) for 30 sec.

Table 1.

Composition of the tablets.		
	Materials	%
1	Citalopram (base)	16.00
2	Kollidon VA64	2.32
3	Lactose 350 mesh	38.98
4	Corn starch	20.00
5	Purified water	25
6	Avicel PH 200 (Microcrystalline cellulose)	20.00
7	Ac-Di-Sol (Croscarmellose sodium)	2.00
8	Magnesium stearate	0.7

[0067] Tablets were produced on a single punch tableting machine Korsch EK0. The characteristics of the tables are shown in Table 2.

Table 2.

Tablet characteristics.	
Parameter	Values
Tablet strength, mg	20
Nominal tablet weight, mg	125
Tablet diameter, mm	7
Tablet shape	Film coating (special doomed)
Mean disintegration time, min	1.77
Mean chcrushing strength, N	69.1
Mean tablet weight, mg	125.4

EP 1 227 088 B1

Table 2. (continued)

Tablet characteristics.	
Parameter	Values
RSD tablet weight, %	0.42
Friability, %	0.3

The tablets produced had satisfactory technical properties.

Example 4

Melt granulation

[0068] The batch size was 200 g. Citalopram base was sieved through a sieve aperture of 0.3 mm. The granulation was performed in a small-scale laboratory high shear mixer (Micromixer)

[0069] The ingredients of the intra-granular phase (1 - 3 in Table 3) were mixed at 1200 rpm.

The jacket temperature was 80 °C. The granulation process was terminated after 3.5 min. The granulate was sieved through a sieve aperture of 1.0 mm and mixed with the extra-granular phase (4, 5) for 3 min. and with the lubricant (6) for 30 sec.

Table 3.

Composition of the tablet.		
	Materials	%
1	Citalopram (base)	16.00
2	Polyethyleneglycol 6000	9.14
3	Lactose 350 mesh	38.98
4	Avicel PH 200 (Microcrystalline cellulose)	30.00
5	Kollidon CL (Cross-linked povidone)	4.00
6	Magnesium stearate	0.7

[0070] Tablets were produced on a single punch tableting machine Korsch EK0. The characteristics of the tables are shown in Table 4.


Table 4.

Tablet characteristics.	
Parameter	Values
Tablet strength, 20 mg	20
Nominal tablet weight, mg	125
Tablet diameter, mm	7
Tablet shape	Film coating, Special doomed
Mean disintegration time, min	1.0
Mean crushing strength, N	55.5
Mean tablet weight, mg	125.6
RSD tablet weight, %	0.5
Friability, %	0.4

[0071] The tablets produced had satisfactory technical properties.

Claims

1. Crystalline base of citalopram, characterised in that it has a purity of more than 99.8 %w/w, preferably more than 99.9 % w/w.
2. Crystalline base of citalopram, prepared by a process characterised in that the base of citalopram is set free and precipitated in crystalline form and optionally re-crystallised one or more times.
3. The crystalline base of citalopram according to claim 2 characterised in that the base of citalopram is set free from a crude salt or a crude mixture of citalopram.
4. Crystalline base of citalopram, prepared by a process which is characterised in that one or more impurities of the formula



wherein Z is halogen, $-O-SO_2-(CF_2)_n-CF_3$, where n is 0-8, $-CHO$, $-NHR^1$, $-COOR^2$, $-CONR^2R^3$ wherein R^2 and R^3 are selected from hydrogen, alkyl, optionally substituted aryl or aralkyl and R^1 is hydrogen or alkylcarbonyl, are removed from a crude mixture of citalopram or from a crude salt of citalopram, by precipitating citalopram base in crystalline form and optionally re-crystallising said base one or more times.


5. The crystalline base of citalopram according to claim 4 wherein the crude mixture of citalopram containing the compound of formula II as an impurity, is prepared by subjecting a compound of formula II to a cyanide exchange reaction with a cyanide source.
6. The crystalline base of citalopram according to claim 4 wherein Z is halogen, in particular bromide or chloride.
7. The crystalline base of citalopram according to any one of claims 4 to 6 wherein the crude mixture of citalopram is subjected to initial purification before the base of citalopram is precipitated in crystalline form.
8. The crystalline base of citalopram according to any one of claims 4 to 6 wherein the crude mixture of citalopram is subjected to initial purification before a crude salt is formed from said crude mixture.
9. The crystalline base of citalopram according to any one of claims 4 to 7 wherein the base of citalopram is set free from a crude salt or a crude mixture of citalopram by treatment with a base and optionally subjected to further purification before the base of citalopram is precipitated in crystalline form.
10. The crystalline base of citalopram according to any one of claims 2 to 4, characterised in that the crude salt is a hydrobromide, hydrochloride, sulphate, oxalate, phosphate or nitrate salt, preferably the sulphate, hydrobromide, or hydrochloride salt.
11. The crystalline base of citalopram according to any one of claims 2 to 10, characterised in that it has a purity of more than 99.8 %w/w, preferably more than 99.9 %w/w.
12. A pharmaceutical composition containing the crystalline base of citalopram according to any one of claims 1 to 11.
13. The pharmaceutical composition according to claim 12 which is a tablet prepared by

- a) direct compression of citalopram, optionally in admixture with pharmaceutically acceptable adjuvants;
 b) compression of a wet granulate of the citalopram, optionally in admixture with pharmaceutically acceptable adjuvants; or
 c) compression of a melt granulate of the citalopram, optionally in admixture with pharmaceutically acceptable adjuvants.

14. The pharmaceutical composition according to claim 12 or 13, characterized in that it contains the racemic mixture of citalopram base.

Patentansprüche

1. Kristalline Base von Citalopram, dadurch gekennzeichnet, daß sie eine Reinheit von mehr als 99,8 % G/G, bevorzugt von mehr als 99,9 % G/G hat.
2. Kristalline Base von Citalopram, hergestellt durch ein Verfahren, das dadurch gekennzeichnet ist, daß die Base von Citalopram freigesetzt und in kristalliner Form ausgefällt und ggf. ein- oder mehrmals umkristallisiert wird.
3. Kristalline Base von Citalopram gemäß Anspruch 2, dadurch gekennzeichnet, daß die Base von Citalopram aus einem rohen Salz oder einer rohen Mischung von Citalopram freigesetzt wird.
4. Kristalline Base von Citalopram, hergestellt durch ein Verfahren, das dadurch gekennzeichnet ist, daß eine oder mehrere Verunreinigungen der Formel



(II)

worin Z Halogen, $-O-SO_2-(CF_2)_n-CF_3$, worin n 0-8 ist, $-CHO$, $-NHR^1$, $-COOR^2$, $-CONR^2R^3$ ist, worin R^2 und R^3 aus Wasserstoff, Alkyl, ggf. substituiertem Aryl oder Aralkyl ausgewählt sind und R^1 Wasserstoff oder Alkylcarbonyl ist, aus einer rohen Mischung von Citalopram oder aus einem rohen Salz von Citalopram entfernt werden, indem Citalopram-Base in kristalliner Form ausgefällt und die Base ggf. ein- oder mehrmals umkristallisiert wird.

5. Kristalline Base von Citalopram gemäß Anspruch 4, worin die rohe Mischung von Citalopram, die die Verbindung der Formel (II) als Verunreinigung enthält, hergestellt wird durch Unterwerfen einer Verbindung der Formel (II) einer Cyanidaustauschreaktion mit einer Cyanidquelle.
6. Kristalline Base von Citalopram gemäß Anspruch 4, worin Z Halogen ist, insbesondere Bromid oder Chlorid.
7. Kristalline Base von Citalopram gemäß einem der Ansprüche 4 bis 6, worin die rohe Mischung von Citalopram einer anfänglichen Reinigung unterworfen wird, bevor die Base von Citalopram in kristalliner Form ausgefällt wird.
8. Kristalline Base von Citalopram gemäß einem der Ansprüche 4 bis 6, worin die rohe Mischung von Citalopram einer anfänglichen Reinigung unterworfen wird, bevor ein rohes Salz aus der rohen Mischung gebildet wird.
9. Kristalline Base von Citalopram gemäß einem der Ansprüche 4 bis 7, worin die Base von Citalopram aus einem rohen Salz oder einer rohen Mischung von Citalopram durch Behandlung mit einer Base freigesetzt und ggf. einer

weiteren Reinigung unterworfen wird, bevor die Base von Citalopram in kristalliner Form ausgefällt wird.

10. Kristalline Base von Citalopram gemäß einem der Ansprüche 2 bis 4, **dadurch gekennzeichnet, daß** das rohe Salz ein Hydrobromid-, Hydrochlorid-, Sulfat-, Oxalat-, Phosphat- oder Nitratsalz ist, bevorzugt das Sulfat-, Hydrobromid- oder Hydrochloridsalz.

11. Kristalline Base gemäß einem der Ansprüche 2 bis 10, **dadurch gekennzeichnet, daß** sie eine Reinheit von mehr als 99,8 % G/G, bevorzugt von mehr als 99,9 % G/G hat.

12. Pharmazeutische Zusammensetzung, die die kristalline Base von Citalopram gemäß einem der Ansprüche 1 bis 11 enthält.


13. Pharmazeutische Zusammensetzung gemäß Anspruch 12, die eine Tablette ist, hergestellt durch

- Direktverpressen von Citalopram, ggf. im Gemisch mit pharmazeutisch akzeptablen Hilfsstoffen;
- Verpressen eines Naßgranulats des Citaloprams, ggf. im Gemisch mit pharmazeutisch akzeptablen Hilfsstoffen; oder
- Verpressen eines Schmelzgranulats des Citaloprams, ggf. im Gemisch mit pharmazeutisch akzeptablen Hilfsstoffen.

14. Pharmazeutische Zusammensetzung gemäß Ansprüchen 12 oder 13, **dadurch gekennzeichnet, daß** sie die racemische Mischung von Citalopram-Base enthält.

Revendications

- Base cristalline de citalopram, **caractérisée en ce qu'elle a une pureté supérieure à 99,8 % en poids, de préférence supérieure à 99,9 % en poids.**
- Base cristalline de citalopram, préparée par un procédé **caractérisé en ce que l'on libère la forme base du citalopram et on la précipite sous forme cristalline, et on la recristallise éventuellement une ou plusieurs fois.**
- Base cristalline de citalopram selon la revendication 2, **caractérisée en ce que l'on libère la forme base du citalopram à partir d'un sel brut ou d'un mélange brut de citalopram.**
- Base cristalline de citalopram, préparée par un procédé **caractérisé en ce que l'on élimine d'un mélange brut de citalopram ou d'un sel brut de citalopram, une ou plusieurs impuretés de formule :**




(II)

dans laquelle Z est un groupe halogéno, $-O-SO_2-(CF_2)_n-CF_3$ dans lequel n est un nombre de 0 à 8, $-CHO$, $-NHR^1$, $-COOR^2$, $-CONR^2R^3$, où R^2 et R^3 sont choisis parmi l'hydrogène, un groupe alkyle, un groupe aryle ou aralkyle éventuellement substitué, et R^1 est un hydrogène ou un groupe alkylcarbonyle, en précipitant le citalopram base sous forme cristalline, et éventuellement en recristallisant cette base une ou plusieurs fois.

5. Base cristalline de citalopram selon la revendication 4, dans laquelle on prépare le mélange brut de citalopram

contenant le composé de formule II comme impureté en soumettant un composé de formule II à une réaction d'échange de cyanure avec une source de cyanure.

- 5 6. Base cristalline de citalopram selon la revendication 4, dans laquelle Z est un halogène, en particulier le chlore ou le brome.
7. Base cristalline de citalopram selon l'une des revendications 4 à 6, dans laquelle on soumet le mélange brut de citalopram à une purification initiale avant de précipiter le citalopram base sous forme cristalline.
- 10 8. Base cristalline de citalopram selon l'une des revendications 4 à 6, dans laquelle on soumet le mélange brut de citalopram à une purification initiale avant de former un sel brut à partir dudit mélange brut.
- 15 9. Base cristalline de citalopram selon l'une des revendications 4 à 7, dans laquelle on libère le citalopram base à partir d'un sel brut ou d'un mélange brut de citalopram par traitement avec une base et on le soumet éventuellement à une purification supplémentaire avant de précipiter le citalopram base sous forme cristalline.
- 20 10. Base cristalline de citalopram selon l'une quelconque des revendications 2 à 4, caractérisée en ce que le sel brut est le bromhydrate, le chlorhydrate, le sulfate, l'oxalate, le phosphate ou le nitrate, de préférence le sulfate, le bromhydrate ou le chlorhydrate.
- 25 11. Base cristalline de citalopram selon l'une quelconque des revendications 2 à 10, caractérisée en ce qu'elle a une pureté supérieure à 99,8 % en poids, de préférence supérieure à 99,9 % en poids.
12. Composition pharmaceutique contenant la base cristalline de citalopram selon l'une des revendications 1 à 11.
- 30 13. Composition pharmaceutique selon la revendication 12, qui est un comprimé préparé par :
 - a) compression directe de citalopram, éventuellement en mélange avec des adjuvants pharmaceutiquement acceptables;
 - b) compression d'un granulé humide de citalopram, éventuellement en mélange avec des adjuvants pharmaceutiquement acceptables ; ou
 - c) compression d'un granulé fondu de citalopram, éventuellement en mélange avec des adjuvants pharmaceutiquement acceptables.
- 35 14. Composition pharmaceutique selon la revendication 12 ou 13, caractérisée en ce qu'elle contient le mélange racémique de citalopram base.



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) **EP 1 227 088 A1**

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
31.07.2002 Bulletin 2002/31

(51) Int Cl.7: **C07D 307/87, A61K 31/343,
A61P 25/24**

(21) Application number: **02009350.6**

(22) Date of filing: **28.02.2001**

(84) Designated Contracting States:
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE TR**
Designated Extension States:
AL LT LV MK RO SI

(30) Priority: **13.03.2000 DK 200000402
13.04.2000 WOPCT/DK00/00183**

(62) Document number(s) of the earlier application(s) in
accordance with Art. 76 EPC:
01909568.6 / 1 169 314

(71) Applicant: **H.Lundbeck A/S
2500 Valby-Copenhagen (DK)**

(72) Inventors:
• **Petersen, Hans
2720 Vanløse (DK)**
• **Bögesö, Klaus Peter
2970 Hörsholm (DK)**
• **Holm, Per
2720 Vanløse (DK)**

(74) Representative: **HOFFMANN - EITLE
Patent- und Rechtsanwälte
Arabellastrasse 4
81925 München (DE)**

Remarks:

This application was filed on 03 - 05 - 2002 as a
divisional application to the application mentioned
under INID code 62.

(54) **Crystalline base of citalopram and hydrochloride or hydrobromide salt thereof**

(57) The present invention relates to the crystalline
base of the well known antidepressant drug citalopram,
1-[3 -(dimethylamino)propyl]-1-(4-fluorophenyl)-1,3-di-
hydro-5-isobenzofuran-carbonitrile, formulations of said
base, a process for the preparation of purified salts of

citalopram, such as the hydrobromide, using the base,
the salts obtained by said process and formulations con-
taining such salts.

EP 1 227 088 A1

Background of the invention


CN(C)CCCC12C=CC(=C1)OC2C3=CC=C(C=C3)F

[0009] This purification process is particularly useful for removing intermediates which are structurally closely related

to citalopram, in particular compounds which only differ from citalopram by the substituent situated in position 5 on the isobenzofuran ring, and intermediates which have physical/chemical properties which are close to those of citalopram, e.g. the 1-[3-(dimethylamino)propyl]-1-(4-fluorophenyl)-1,3-dihydro-isobenzofuranes having halogen (in particular bromide and chloride), an amide or an ester in position 5 of the isobenzofuran ring, or 1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofuranbromide, or -chloride.

Summary of the invention

[0010] The present invention provides the crystalline base of the compound



(I)


[0011] In a second aspect, the invention provides a process for the manufacture of a salt of citalopram, preferably the hydrobromide or hydrochloride in which the free base of citalopram is precipitated in crystalline form, optionally re-crystallised one or more times and then transferred to a pharmaceutically acceptable salt of citalopram.

[0012] In a further aspect, the invention relates to the pure crystalline salt, preferably the hydrobromide or hydrochloride prepared by the process of the invention.

[0013] In particular, the invention relates to a process for the manufacture of a salt of citalopram characterised in that the base of citalopram is set free and precipitated in crystalline form, optionally re-crystallised one or more times, and then transferred into a salt thereof.

[0014] In particular, the invention relates to a process for the manufacture of a salt of citalopram characterised in that the base of citalopram is set free from a crude salt or crude mixture of citalopram.

[0015] More particularly, the present invention relates to a process for the manufacture of citalopram base or a salt of citalopram characterised in that one or more impurities of the formula



(II)

wherein Z is halogen, $-O-SO_2-(CF_2)_n-CF_3$, where n is 0-8, $-CHO$, $-NHR^1$, $-COOR^2$, $-CONR^2R^3$ wherein R^2 and R^3 are selected from hydrogen, alkyl, optionally substituted aryl or aralkyl and R^1 is hydrogen or alkylcarbonyl, are removed from a crude mixture of citalopram or from a crude salt of citalopram, by precipitating citalopram base in crystalline form, optionally re-crystallising said base one or more times and/or transferring said base into a salt thereof.

[0016] The crude mixture of citalopram containing the compound of formula II as an impurity may be prepared by subjecting a compound of formula II to a cyanide exchange reaction with a cyanide source, or by subjecting 1-(4-fluorophenyl)-1,3-dihydro-5-isobenzofuranhalogenide, in particular the bromide, to a cyanide exchange reaction followed by alkylation with a 3-(N,N-dimethylamino)propyl-halogenide.

[0017] In a particular embodiment of the invention, Z is halogen, in particular bromide or chloride.

[0018] In a particularly preferred embodiment of the invention, the salt prepared is the hydrobromide or hydrochloride salt of citalopram.

[0019] The crude salt may be any convenient salt, such as the hydrobromide, hydrochloride, sulphate, oxalate, phosphate, nitrate or any other convenient salt. Other salts are salts of organic acids.

[0020] In a preferred embodiment of the invention, the crude salt is the sulphate, the hydrobromide or the hydrochloride salt.

5 [0021] The invention also relates to a hydrochloride or hydrobromide salt of citalopram prepared by the processes of the invention. In particular, the invention relates to a hydrochloride or hydrobromide salt of citalopram having a purity of more than 99.8 %w/w, preferably more than 99.9 %w/w.

[0022] In yet another aspect, a pharmaceutical formulation of the free base of citalopram, or a hydrobromide or hydrochloride prepared from said base, is provided. Preferably the formulation is for oral administration.

10 [0023] The formulations according to the invention may be prepared by direct compression of citalopram in admixture with conventional adjuvants or diluents. Alternatively, a wet granulate or a melt granulate of citalopram, optionally in admixture with conventional adjuvants or diluents may be used for compression of tablets.

[0024] In particular, the pharmaceutical composition of the invention contains the racemic mixture of citalopram base, citalopram hydrochloride or citalopram hydrobromide.

15 [0025] The crystalline base of citalopram is preferably more than 99.8% w/w pure, most preferably more than 99.9% w/w pure (peak area). The melting point is preferably a range within 90 - 93 °C, most preferably 91 - 92 °C (DSC; onset, open capsule) or it is between 92 and 94°C, preferably 92.5 and 93.5 °C (DSC; onset, closed capsule). The crystalline base of citalopram is preferably in racemic form.

[0026] The terms "crude salt" and "crude mixture" refer to the fact that the salt and the mixture, respectively, comprise impurities, in particular impurities of formula II, which must be removed or which it is desired to remove.

20 [0027] The crude salt may be a salt separated directly from the reaction mixture, or the crude reaction mixture may have been subjected to some initial purification, e.g. one re-crystallisation, and /or treatment with activated carbon or silica gel, and the salt formed subsequently by treatment with an acid using methods known in the art. The salt may be isolated by precipitation or it may exist in a solvent, e.g. in the mixture resulting directly from the synthesis of the salt.

25 [0028] Similarly, the crude mixture comprising citalopram may be obtained directly from the synthesis of the compound according to any of the above mentioned processes or it may have been subjected to some initial or simultaneous purification, e.g. one re-crystallisation, treatment with activated carbon or silica gel.

[0029] The base of citalopram may be set free from the crude salt by dissolving the crude salt in a mixture of water and an organic solvent and then adding a base. The organic solvent may be toluene, ethyl acetate or any other suitable solvent and the base may be any convenient base, preferably NaOH or NH₃. Likewise, the base of citalopram may, if necessary, be set free from a crude mixture containing citalopram by treatment with a base.

30 [0030] Crude mixtures containing citalopram base may be subjected to further purification and extraction, before the base is precipitated in crystalline form. The base of citalopram may be isolated by separation of the organic phase, evaporation of the solvent in order to obtain the base most probably as an oil and then crystallisation of the base from an aprotic solvent, such as an alkane, including n-heptane, hexane and isooctane, and high and low boiling petroleum ethers and substituted aromates, inc1 toluene and xylenes. Crystalline citalopram base may be re-crystallised from the same solvents.

35 [0031] The pharmaceutically acceptable salt of citalopram, such as the hydrobromide or hydrochloride, may be prepared by methods known in the art. So, the base may be reacted with either the calculated amount of acid in a water miscible solvent, such as acetone or ethanol, with subsequent isolation of the salt by concentration and cooling, or with an excess of the acid in a water immiscible solvent, such as ethylether, ethylacetate or dichloromethane, with the salt separating spontaneously. The hydrobromide or hydrochloride of citalopram obtained by the method of the invention has a very high purity, preferably more than 99.8% pure, most preferably more than 99.9 % purity. Other salts of citalopram, e.g. the oxalate, may also be obtained in a very pure form by this process.

40 [0032] The cyanide exchange reactions mentioned above may be carried out as described in the patent applications mentioned above.

[0033] In particular, when Z is halogen, or CF₃-(CF₂)_n-SO₂-O- wherein n is an integer in the range 0-8, incl., the conversion to a cyano group may be carried out by reaction with a cyanide source, for example KCN, NaCN, CuCN, Zn(CN)₂ or (R⁴)₄NCN where R⁴ indicates four groups which may be the same or different and are selected from hydrogen and straight chain or branched alkyl, in the presence of a palladium catalyst and a catalytic amount of Cu⁺ or Zn²⁺, or with Zn(CN)₂ in the presence of a palladium catalyst.

45 [0034] The cyanide source is used in a stoichiometric amount or in excess, preferably 1-2 equivalents are used pr. equivalent starting material. R⁴N⁺ may conveniently be (Bu)₄N⁺. The cyanide compound is preferably NaCN or KCN or Zn(CN)₂.


50 [0035] The palladium catalyst may be any suitable Pd(0) or Pd(II) containing catalyst, such as Pd(PPh₃)₄, Pd₂(dba)₃, Pd(PPh₃)₂Cl₂, etc. The Pd catalyst is conveniently used in an amount of 1-10, preferably 2-6, most preferably about 4-5 mol%.

[0036] Catalytic amounts of Cu⁺ and Zn²⁺, respectively, means substoichiometric amounts such as 0.1 - 5, preferably

1 - 3 %. Conveniently, about ½ eq. is used per eq. Pd. Any convenient source of Cu^+ and Zn^{2+} may be used. Cu^+ is preferably used in the form of CuI and Zn^{2+} is conveniently used as the $\text{Zn}(\text{CN})_2$ salt.

[0037] When Z is Br or I, the conversion to a cyano group may also be carried out by reaction with $\text{Cu}(\text{CN})$ without catalyst. In a preferred embodiment, the reaction is performed at elevated temperature.

[0038] In another aspect of the invention, the reaction is performed in an ionic liquid of the general formula $(\text{R}^5)_4\text{N}^+$, X^- , wherein R^5 are alkyl-groups or two of the R^5 groups together form a ring and X^- is the counterion. In one embodiment of the invention, $(\text{R}^5)_4\text{N}^+\text{X}^-$ represents



[0039] In another particular aspect, the reaction is conducted with apolar solvents such as benzene, xylene or mesitylene and under the influence of microwaves by using *i.e.* Synthrowave 1000™ by Prolabo. In a particular aspect, the reaction is performed without added solvent.

[0040] The temperature ranges are dependent upon the reaction type. If no catalyst is present, preferred temperatures are in the range of 100-200°C. However, when the reaction is conducted under the influence of microwaves, the temperature in the reaction mixture may raise to above 300 °C. More preferred temperature ranges are between 120-170°C. The most preferred range is 130-150°C. If catalyst is present, the preferred temperature range is between 0 and 100°C. More preferred are temperature ranges of 40-90°C. Most preferred temperature ranges are between 60-90°C.

[0041] Other reaction conditions, solvents, etc. are conventional conditions for such reactions and may easily be determined by a person skilled in the art.

[0042] When Z is Cl or Br, the conversion to a cyano group may also be carried out by reaction with a cyanide source, for example KCN , NaCN , CuCN , $\text{Zn}(\text{CN})_2$ or $(\text{R}^4)_4\text{NCN}$ where $(\text{R}^4)_4$ indicates four groups which may be the same or different and are selected from hydrogen and straight chain or branched alkyl, in the presence of a nickel catalyst.

[0043] The nickel catalyst may be any suitable $\text{Ni}(0)$ or $\text{Ni}(\text{II})$ containing complex which acts as a catalyst, such as $\text{Ni}(\text{PPh}_3)_3$, $(\sigma\text{-aryl})\text{-Ni}(\text{PPh}_3)_2\text{Cl}$, etc. The nickel catalysts and their preparation are described in WO 96/11906, EP-A-613720 or EP-A-384392.

[0044] In one embodiment of the invention, the reaction is carried out in the presence of a catalytic amount of Cu^+ or Zn^{2+} .

[0045] In a particularly preferred embodiment, a Nickel(0) complex is prepared *in situ* before the cyanation reaction by reduction of a Nickel(II) precursor such as NiCl_2 or NiBr_2 by a metal, such as zinc, magnesium or mangan in the presence of excess of complex ligands, preferably triphenylphosphine.

[0046] The Ni-catalyst is conveniently used in an amount of 0.5-10, preferably 2-6, most preferably about 4-5 mol%.

[0047] Catalytic amounts of Cu^+ and Zn^{2+} , respectively, mean substoichiometric amounts such as 0.1 - 5, preferably 1 - 3 %. Any convenient source of Cu^+ and Zn^{2+} may be used. Cu^+ is preferably used in the form of CuI and Zn^{2+} is conveniently used as the $\text{Zn}(\text{CN})_2$ salt or formed *in situ* by reduction of a Nickel (II) compounds using zinc.

[0048] The Ni catalysts are *i.e.* $\text{Ni}(0)$, $\text{Pd}(0)$ or $\text{Pd}(\text{II})$ catalysts as described by Sakakibara *et. al.* in Bull. Chem. Soc. Jpn., 61, 1985-1990, (1988). Preferred catalysts are $\text{Ni}(\text{PPh}_3)_3$ or $\text{Pd}(\text{PPh}_3)_4$, or $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$.

[0049] The reactions may be performed in any convenient solvents described in Sakakibara *et. al.* in Bull. Chem. Soc. Jpn., 61, 1985-1990, (1988). Preferred solvents are acetonitril, ethylacetat, THF, DMF or NMP.

[0050] When Z is CHO, the conversion to a cyano group may be carried out by conversion of the formyl group to an oxime or similar group by reaction with a reagent $\text{R}^6\text{-V-NH}_2$ wherein R^6 is hydrogen, optionally substituted alkyl, aryl or heteroaryl and V is O, N or S, followed by dehydration with a common dehydrating agent, for example thionylchloride, acetic anhydride/pyridine, pyridine/HCl or phosphor pentachloride. Preferred reagents $\text{R}^6\text{-V-NH}_2$ are hydroxylamin and compounds wherein R^6 is alkyl or aryl and V is N or O.

[0051] When Z is $-\text{COOH}$, the conversion to a cyano group may be carried out via the corresponding acid chloride, ester or amide.

[0052] The acid chloride is conveniently obtained by treatment of the acid with POCl_3 , PCl_5 or SOCl_2 neat or in a suitable solvent, such as toluene or toluene comprising a catalytic amount of N,N-dimethylformamide. The ester is obtained by treatment of the acid with an alcohol, in the presence of an acid, preferably a mineral acid or a Lewis acid, such as HCl, H_2SO_4 , POCl_3 , PCl_5 or SOCl_2 . Alternatively, the ester may be obtained from the acid chloride by reaction with an alcohol. The ester or the acid chloride is then converted to an amide or by amidation with ammonia or an

alkylamine, preferably t-butyl amine.

[0053] The conversion to amide may also be obtained by reaction of the ester with ammonia or an alkylamine under pressure and heating.

[0054] The amide group is then converted to a cyano group by dehydration. The dehydrating agent may be any suitable dehydrating agent, and the optimal agent may easily be determined by a person skilled in the art. Examples of suitable dehydrating agents are SOCl_2 , POCl_3 and PCl_5 , preferably SOCl_2 .

[0055] In a particularly preferred embodiment, the carboxylic acid is reacted with an alcohol, preferably ethanol, in the presence of POCl_3 , in order to obtain the corresponding ester, which is then reacted with ammonia thereby giving the corresponding amide, which in turn is reacted with SOCl_2 in toluene comprising a catalytic amount of N,N-dimethylformamide.

[0056] Alternatively, a compound where Z is $-\text{COOH}$ may be reacted with chlorosulfonyl isocyanate in order to form the nitrile, or treated with a dehydrating agent and a sulfonamide.

[0057] When Z is $-\text{NHR}^1$, where R^1 is hydrogen, the conversion into cyano is preferably performed by diazotation and followed by reaction with CN^- . Most preferably NaNO_2 and CuCN and/or NaCN are used. When R^1 is alkylcarbonyl, it is initially subjected to hydrolysis thereby obtaining the corresponding compound wherein R^1 is H which is the converted as described above. The hydrolysis may be performed either in acidic or basic environment.

[0058] The compounds of formula (II) may be prepared as described in DE 2,657,013, WO 0011926 and WO 0013648, WO 9819513, WO 9819512 and WO 9900548.

[0059] Throughout this specification with claims halogen means chloro, bromo or iodo.

[0060] The term alkyl refers to a branched or unbranched alkyl group, such as methyl, ethyl, 1-propyl, 2-propyl, 1-butyl, 2-butyl, 2-methyl-2-propyl, and 2-methyl-1-propyl.

[0061] The term aryl refers to a carbocyclic aromatic group, in particular phenyl. Aralkyl refers to an arylalkyl group wherein aryl and alkyl is as defined above. The aryl and aralkyl groups may optionally be substituted, e.g. with alkyl groups, forming for example tolyl.

[0062] The pharmaceutical compositions of the invention may be administered in any suitable way and in any suitable form, for example orally in the form of tablets, capsules, powders or syrups, or parenterally in the form of usual sterile solutions for injection. Preferably the pharmaceutical compositions of the invention are administered orally.

[0063] The pharmaceutical formulations of the invention may be prepared by conventional methods in the art. For example, tablets may be prepared by mixing the active ingredient with ordinary adjuvants and/or diluents and subsequently compressing the mixture in a conventional tableting machine. Examples of adjuvants or diluents comprise: Corn starch, potato starch, talcum, magnesium stearate, gelatine, lactose, gums, and the like. Any other adjuvant or additive colourings, aroma, preservatives etc. may be used provided that they are compatible with the active ingredients.

[0064] In particular, the formulations according to the invention may be prepared by direct compression of citalopram in admixture with conventional adjuvants or diluents. Alternatively, a wet granulate or a melt granulate of citalopram, optionally in admixture with conventional adjuvants or diluents may be used for compression of tablets.

[0065] Solutions for injections may be prepared by solving the active ingredient and possible additives in a part of the solvent for injection, preferably sterile water, adjusting the solution to the desired volume, sterilisation of the solution and filling in suitable ampoules or vials. Any suitable additive conventionally used in the art may be added, such as tonicity agents, preservatives, antioxidants, etc.

[0066] According to the present invention, the base of citalopram has been found to be crystalline with stable and nice white crystals and it has been found that the base may easily be crystallised in a very pure form. So for example more than 99.8% w/w pure citalopram base was obtained by crystallisation from up to 95% pure hydrobromide without further purification. Accordingly, the process of the invention for preparing salts of citalopram has been found to give the salts as very pure products of pharmaceutically acceptable quality. Accordingly, the yield may be improved substantially during the manufacture of citalopram.

[0067] Finally, it has been found that the crystalline citalopram base may be formulated into very good and stable solid formulations with good release properties.

[0068] The invention is further illustrated by the following examples.

Example 1

Crystallisation of R,S-Citalopram as the free base.

[0069] 1-(3-Dimethylaminopropyl)-1-(4'-fluorophenyl)-1,3-dihydrobenzofuran-5-carbonitrile.

[0070] 1-(3-Dimethylaminopropyl)-1-(4'-fluorophenyl)-1,3-dihydrobenzofuran-5-carbonitrile hydrobromide (101 grams, 0.25 mole) prepared from 1-(3-Dimethylaminopropyl)-1-(4'-fluorophenyl)-1,3-dihydro-5-isobenzofuranbromide, is suspended in water (500 ml) and toluene (500 ml). NaOH (60 ml, 5 N (aq)) is added and the mixture ($\text{pH} > 10$) is

stirred for 15 min. before the phases are separated. The organic phase is washed with water (2x100 ml) and filtered through a pad of filter help. The volatiles are removed *in vacuo* and the title compound is obtained as an oil. n-Heptane (400 ml) is added and the mixture is heated to 70 °C. On cooling, crystals form. The white crystals of the title compound are filtered off and dried at ambient temperature over night *in vacuo*. Yield: 75.4 grams (93%). DSC(onset, open capsule): 91.3-91.8 °C DSC (onset, closed capsule): 92.8 °C. Purity: (> 99.8 % (peak area)).

Anal. calcd. for C₂₀H₂₁N₂F₁₀O₁; C, 74.04; H, 6.54; N, 8.64. Found C, 74.01; H, 6.49; N, 8.59. ¹H-NMR (DMSO-d₆, 500 MHz): 1.21 (1H, m), 1.29 (1H, m), 2.02 (6H, s), 2.09-2.23 (4 H, m), 5.15 (1H, d J=12.5 Hz), 5.22 (1H, d J=12.5 Hz), 7.16 (2H, t J=8.5 Hz), 7.60 (2H, dt J=8.5 Hz J=1.2 Hz), 7.76 (1H, d J= 8.5 Hz), 7.79 (1H, d J=8.5 Hz), 7.80 (1H, s). ¹³C-NMR (DMSO-d₆, 125 MHz): 21.8, 38.3, 45.0, 58.8, 71.0, 90.7, 110.5, 115.1 (d J=22 Hz), 118.8, 123.1, 125.1, 127.0 (d J=8 Hz), 132.0, 140.0 (d J=3 Hz), 140.5, 149.5, 161.3 (d J=245Hz).

Example 2

[0071]

a) A crude mixture of Citalopram and sulphuric acid is made basic by adding NaOH and the citalopram base is extracted with toluene. The toluene is evaporated and the citalopram base obtained is dissolved in n-heptane at elevated temperature. The very pure free base of citalopram is precipitated by cooling.

b) A crude mixture of Citalopram and sulphuric acid is made basic by adding NaOH and the citalopram base is extracted with toluene. The toluene is evaporated and the citalopram base obtained is dissolved in methanol. The mixture is treated with activated carbon and filtrated and the solvent is evaporated. The purified free base is dissolved in n-heptane at elevated temperature. Then the very pure free base of citalopram is precipitated by cooling.

c) A crude mixture of Citalopram and sulphuric acid is made basic by adding NaOH and the citalopram base is extracted with toluene. The toluene phase is treated with silicagel, the toluene is evaporated and the citalopram base obtained is dissolved in n-heptane at elevated temperature. The very pure free base of citalopram is precipitated by cooling.

d) A crude mixture of Citalopram and sulphuric acid is made basic by adding NaOH and the citalopram base is extracted with toluene. The toluene phase is treated with silicagel, the toluene is evaporated and the citalopram base obtained is dissolved in methanol. The mixture is treated with activated carbon and filtrated and the solvent is evaporated. The purified free base is dissolved in n-heptane at elevated temperature. Then the extremely pure free base of citalopram is precipitated by cooling.

Example 3

Wet granulation and preparation of tablets

[0072] The batch size was 200 g and the granulation was performed in a small-scale laboratory high shear mixer (Micromixer).

[0073] Citalopram base was sieved through a sieve aperture of 0.3 mm. The ingredients of the intragranular phase (1 - 4 in Table 1) were mixed at 600 rpm. 25 ml of purified water (5) was added in 30 sec and the granulation terminated after a total processing time of 3 min. The granulate was wet sieved through a 0.7 mm sieve aperture and dried at 40 °C in 30 minutes to equilibrium relative humidity of 32 %. The dried granulate was finally sieved through a 0.7 mm sieve aperture.

[0074] The dried granulate was mixed for 3 minutes with the extragranular phase (6- 7) in a Turbula mixer and finally mixed with the lubricant (8) for 30 sec.

Table 1.

Composition of the tablets.		
	Materials	%
1	Citalopram (base)	16.00
2	Kollidon VA64	2.32
3	Lactose 350 mesh	38.98

Table 1. (continued)

Composition of the tablets.		
	Materials	%
4	Corn starch	20.00
5	Purified water	25
6	Avicel PH 200 (Microcrystalline cellulose)	20.00
7	Ac-Di-Sol (Croscarmellose sodium)	2.00
8	Magnesium stearate	0.7

[0075] Tablets were produced on a single punch tableting machine Korsch EK0. The characteristics of the tables are shown in Table 2.

Table 2.

Tablet characteristics.	
Parameter	Values
Tablet strength, mg	20
Nominal tablet weight, mg	125
Tablet diameter, mm	7
Tablet shape	Film coating (special doomed)
Mean disintegration time, min	1.77
Mean chrushing strength, N	69.1
Mean tablet weight, mg	125.4
RSD tablet weight, %	0.42
Friability, %	0.3

[0076] The tablets produced had satisfactory technical properties.

Example 4

Melt granulation

[0077] The batch size was 200 g. Citalopram base was sieved through a sieve aperture of 0.3 mm. The granulation was performed in a small-scale laboratory high shear mixer (Micromixer)

[0078] The ingredients of the intra-granular phase (1 - 3 in Table 3) were mixed at 1200 rpm.

The jacket temperature was 80 °C. The granulation process was terminated after 3.5 min. The granulate was sieved through a sieve aperture of 1.0 mm and mixed with the extra-granular phase (4, 5) for 3 min. and with the lubricant (6) for 30 sec.

Table 3.

Composition of the tablet.		
	Materials	%
1	Citalopram (base)	16.00
2	Polyethyleneglycol 6000	9.14
3	Lactose 350 mesh	38.98
4	Avicel PH 200 (Microcrystalline cellulose)	30.00
5	Kollidon CL (Cross-linked povidone)	4.00

Table 3. (continued)

Composition of the tablet.		
	Materials	%
6	Magnesium stearate	0.7

[0079] Tablets were produced on a single punch tableting machine Korsch EK0. The characteristics of the tables are shown in Table 4.


Table 4.

Tablet characteristics.	
Parameter	Values
Tablet strength, 20 mg	20
Nominal tablet weight, mg	125
Tablet diameter, mm	7
Tablet shape	Film coating , Special doomed
Mean disintegration time, min	1.0
Mean chrushing strength, N	55.5
Mean tablet weight, mg	125.6
RSD tablet weight, %	0.5
Friability, %	0.4

[0080] The tablets produced had satisfactory technical properties.

Claims

1. A crystalline base of citalopram, or a hydrochloride or hydrobromide salt of citalopram, **characterised in that** it has a purity of more than 99.8 %w/w, preferably more than 99.9 % w/w.
2. The crystalline base of citalopram, or a hydrochloride or hydrobromide salt of citalopram prepared by a process **characterised in that** the base of citalopram is set free and precipitated in crystalline form, optionally re-crystallised one or more times, and then transferred into a salt thereof.
3. The crystalline base of citalopram, or a hydrochloride or hydrobromide salt of citalopram of claim 2 **characterised in that** the base of citalopram is set free from a crude salt or a crude mixture of citalopram.
4. The crystalline base of citalopram, or a hydrochloride or hydrobromide salt of citalopram prepared by a process **characterised in that** one or more impurities of the formula



(II)

wherein Z is halogen, $-O-SO_2-(CF_2)_n-CF_3$, where n is 0-8, $-CHO$, $-NHR^1$, $-COOR^2$, $-CONR^2R^3$ wherein R^2 and R^3 are selected from hydrogen, alkyl, optionally substituted aryl or aralkyl and R^1 is hydrogen or alkylcarbonyl, are removed from a crude mixture of citalopram or from a crude salt of citalopram, by precipitating citalopram base in crystalline form, optionally re-crystallising said base one or more times and/or transferring said base into a salt thereof.

5 5. The crystalline base of citalopram, or a hydrochloride or hydrobromide salt of citalopram according to claim 4 wherein the crude mixture of citalopram containing the compound of formula II as an impurity, is prepared by subjecting a compound of formula II to a cyanide exchange reaction with a cyanide source.

10 6. The crystalline base of citalopram, or a hydrochloride or hydrobromide salt of citalopram of claim 4 wherein Z is halogen, in particular bromide or chloride.

15 7. The crystalline base of citalopram, or a hydrochloride or hydrobromide salt of citalopram of claims 4-6 wherein the crude mixture of citalopram is subjected to initial purification before the base of citalopram is precipitated in crystalline form.

20 8. The crystalline base of citalopram, or a hydrochloride or hydrobromide salt of citalopram of claims 4-6 wherein the crude mixture of citalopram is subjected to initial purification before a crude salt is formed from said crude mixture.

9. The crystalline base of citalopram, or a hydrochloride or hydrobromide salt of citalopram according to claims 4-7 wherein the base of citalopram is set free from a crude salt or a crude mixture of citalopram by treatment with a base and optionally subjected to further purification before the base of citalopram is precipitated in crystalline form.

25 10. The crystalline base of citalopram, or a hydrochloride or hydrobromide salt of citalopram according to claims 2-9 wherein the citalopram base is transferred into the hydrobromide or the hydrochloride salt of citalopram.

30 11. The crystalline base of citalopram, or a hydrochloride or hydrobromide salt of citalopram according to claims 2-4, characterised in that the crude salt is a hydrobromide, hydrochloride, sulphate, oxalate, phosphate or nitrate salt, preferably the sulphate hydrobromide, or hydrochloride salt.

12. The base, the hydrochloride or the hydrobromide salt of claims 2-11 characterised in that it has a purity of more than 99.8 %w/w, preferably more than 99.9 %w/w.

35 13. A pharmaceutical composition containing the hydrochloride or the hydrobromide salt of citalopram according to Claims 1 to 12, or the crystalline base of citalopram.

14. A pharmaceutical composition according to Claim 13 which is a tablet prepared by

- 40 a) direct compression of citalopram, optionally in admixture with pharmaceutically acceptable adjuvants;
b) by compression of a wet granulate of the citalopram, optionally in admixture with pharmaceutically acceptable adjuvants; or
c) by compression of a melt granulate of the citalopram, optionally in admixture with pharmaceutically acceptable adjuvants.

45 15. The pharmaceutical composition according to Claims 13 to 14 characterized in that it contains the racemic mixture of citalopram base, citalopram hydrochloride or citalopram hydrobromide.



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 02 00 9350

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (InCL7)
X	EP 0 171 943 A (LUNDBECK, H., OG CO. A/S, DEN.) 19 February 1986 (1986-02-19)	1,13-15	C07D307/87 A61K31/343 A61P25/24
A	* page 3 - page 4; example II * * page 7 - page 8; example 2 * * page 8, last paragraph *	2-12	
X	DE 26 57 013 A (KEFALAS A/S, DEN.) 28 July 1977 (1977-07-28)	1,13-15	
A	* page 13, line 6 - line 15; claims * * page 19 - page 21; example 3 *	2-12	
A	EP 0 347 066 A (LUNDBECK, H., OG CO. A/S, DEN.) 20 December 1989 (1989-12-20)	1-15	
A,D	WO 98 19511 A (H. LUNDBECK A/S, DEN.; PETERSEN, HANS; BOGESO, KLAUS PETER; BECH SOMMER) 14 May 1998 (1998-05-14)	1-15	
	* page 8, line 20 - line 35 * * page 8, line 36 - line 37 *		
A,D	WO 98 19512 A (H. LUNDBECK A/S, DEN.; PETERSEN, HANS; BREGNEDAL, PETER; BOGESO, KLAUS) 14 May 1998 (1998-05-14)	1-15	TECHNICAL FIELDS SEARCHED (InCL7) C07D A61K A61P
	* page 7, line 28 - line 38; claim 1 *		
A,D	WO 98 19513 A (H. LUNDBECK A/S, DEN.; PETERSEN, HANS; BREGNEDAL, PETER; BOGESO, KLAUS) 14 May 1998 (1998-05-14)	1-15	
	* page 10 - page 11; claim 1; example 6 *		
P,X	DE 200 07 303 U (H. LUNDBECK A/S, DEN.) 31 August 2000 (2000-08-31)	1-15	
	* the whole document *		
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 6 June 2002	Examiner Paisdor, B
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document</p> <p>T: theory or principle underlying the invention E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document</p>			

EPO FORM 1503 (03.02) (P04031)

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 02 00 9350

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

06-06-2002

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0171943	A	19-02-1986	AT 38661 T	15-12-1988
			AU 574819 B2	14-07-1988
			AU 4577685 A	13-02-1986
			CA 1237147 A1	24-05-1988
			DE 3566251 D1	22-12-1988
			DK 89595 A	10-08-1995
			DK 356285 A	07-02-1986
			EP 0171943 A1	19-02-1986
			ES 545885 D0	01-04-1986
			ES 8606257 A1	01-10-1986
			FI 852902 A ,B,	07-02-1986
			GR 851894 A1	03-12-1985
			IE 57817 B1	21-04-1993
			IL 75690 A	31-10-1988
			JP 1902596 C	08-02-1995
			JP 6025099 B	06-04-1994
			JP 61087654 A	06-05-1986
			NO 853091 A ,B,	07-02-1986
			NZ 212541 A	30-05-1988
			PT 80913 A ,B	01-09-1985
			US 4650884 A	17-03-1987
			ZA 8505026 A	25-06-1986
DE 2657013	A	28-07-1977	GB 1526331 A	27-09-1978
			AT 360001 B	10-12-1980
			AT 571979 A	15-05-1980
			AT 360002 B	10-12-1980
			AT 572079 A	15-05-1980
			AT 359488 B	10-11-1980
			AT 947276 A	15-04-1980
			AU 509445 B2	15-05-1980
			AU 2107377 A	13-07-1978
			BE 850401 A1	14-07-1977
			CA 1094087 A1	20-01-1981
			CH 626886 A5	15-12-1981
			CH 632258 A5	30-09-1982
			CH 632259 A5	30-09-1982
			DE 2657013 A1	28-07-1977
			DK 13177 A ,B,	15-07-1977
			ES 454980 A1	01-04-1978
			FI 770073 A ,B,	15-07-1977
			FR 2338271 A1	12-08-1977
			IE 44055 B1	29-07-1981
			JP 1368581 C	11-03-1987
			JP 52105162 A	03-09-1977
			JP 61035986 B	15-08-1986

EPO FORM P459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 02 00 9350

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

06-06-2002

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
DE 2657013	A		NL 7700244 A ,B,	18-07-1977
			NO 770109 A ,B,	15-07-1977
			NZ 183001 A	02-06-1978
			SE 429551 B	12-09-1983
			SE 7614201 A	15-07-1977
			US 4136193 A	23-01-1979
			ZA 7700057 A	30-11-1977
EP 0347066	A	20-12-1989	AT 119896 T	15-04-1995
			AU 623144 B2	07-05-1992
			AU 3629589 A	04-01-1990
			CA 1339568 A1	02-12-1997
			CY 2081 A	16-10-1998
			DE 68921672 D1	20-04-1995
			DE 68921672 T2	27-07-1995
			DK 11593 A	01-02-1993
			DK 259989 A	15-12-1989
			EP 0347066 A1	20-12-1989
			ES 2068891 T3	01-05-1995
			FI 892823 A ,B,	15-12-1989
			FI 941829 A ,B,	20-04-1994
			FI 20000507 A	06-03-2000
			GR 3015889 T3	31-07-1995
			HK 139596 A	02-08-1996
			HU 9500496 A3	28-09-1995
			IE 65734 B1	15-11-1995
			IL 90465 A	24-01-1995
			JP 3038204 B2	08-05-2000
			JP 11292867 A	26-10-1999
			JP 2036177 A	06-02-1990
			JP 3044253 B2	22-05-2000
			MX 9203346 A1	31-08-1992
			NO 892447 A ,B,	15-12-1989
			NZ 229426 A	21-12-1990
			PT 90845 A ,B	29-12-1989
			US RE34712 E	30-08-1994
			US 4943590 A	24-07-1990
			ZA 8904476 A	25-04-1990
WO 9819511	A	14-05-1998	WO 9819511 A2	14-05-1998
			AU 738526 B2	20-09-2001
			AU 5116798 A	29-05-1998
			BG 104487 A	31-01-2001
			BR 9714924 A	26-09-2000
			DE 1032566 T1	15-03-2001
			EP 1032566 A2	06-09-2000

EPO FORM P4489

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 02 00 9350

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

06-06-2002

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9819511	A		ES 2149141 T1	01-11-2000
			NO 20002267 A	06-07-2000
			PL 340774 A1	26-02-2001
			SK 6812000 A3	09-10-2000
			TR 200001314 T2	21-11-2000
			US 6291689 B1	18-09-2001
			ZA 9810059 A	05-05-1999
WO 9819512	A	14-05-1998	WO 9819512 A2	14-05-1998
			AU 738359 B2	13-09-2001
			AU 5116898 A	29-05-1998
			BG 104486 A	31-01-2001
			DE 1042310 T1	19-04-2001
			EP 1042310 A2	11-10-2000
			ES 2149734 T1	16-11-2000
			HU 0002953 A2	28-04-2001
			NO 20002077 A	10-05-2000
			NZ 504069 A	26-10-2001
			PL 340605 A1	12-02-2001
			SK 6822000 A3	09-10-2000
			TR 200001341 T2	21-11-2000
			US 6258842 B1	10-07-2001
			ZA 9810058 A	05-05-1999
WO 9819513	A	14-05-1998	AT 205824 T	15-10-2001
			AU 737610 B2	23-08-2001
			AU 6609898 A	29-05-1998
			BG 104116 A	29-12-2000
			BR 9810499 A	13-03-2001
			CN 1268129 T	27-09-2000
			DE 69801764 D1	25-10-2001
			DE 1015416 T1	05-10-2000
			WO 9819513 A2	14-05-1998
			DK 1015416 T3	05-11-2001
			EP 1015416 A2	05-07-2000
			ES 2148120 T1	16-10-2000
			HU 0003177 A2	28-04-2001
			JP 2002509526 T	26-03-2002
			NO 20000008 A	03-01-2000
			NZ 501737 A	26-10-2001
			PL 338015 A1	25-09-2000
			SI 1015416 T1	31-12-2001
			SK 32000 A3	11-07-2000
			TR 200000066 T2	21-11-2000
			US 6229026 B1	08-05-2001
			ZA 9805981 A	28-01-1999

EPO FORM P449

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

**ANNEX TO THE EUROPEAN SEARCH REPORT
ON EUROPEAN PATENT APPLICATION NO.**

EP 02 00 9350

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report.
The members are as contained in the European Patent Office EDP file on
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

06-06-2002

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 20007303 U	27-07-2000	AT 4364 U1	25-06-2001
		AU 3725201 A	13-09-2001
		BE 1013210 A3	02-10-2001
		CH 691477 A5	31-07-2001
		CH 691537 A5	15-08-2001
		CZ 20010808 A3	16-01-2002
		DE 10108042 A1	18-10-2001
		DE 20007303 U1	27-07-2000
		WO 0168627 A1	20-09-2001
		DK 173903 B1	11-02-2002
		EP 1169314 A1	09-01-2002
		ES 2159491 A1	01-10-2001
		FI 20010225 A	14-09-2001
		FR 2806086 A1	14-09-2001
		GB 2357762 A ,B	04-07-2001
		GR 1003796 B2	08-02-2002
		NL 1016435 C1	06-11-2000
		NL 1017413 C1	13-09-2001
		NO 20010619 A	14-09-2001
		NO 20020356 A	14-09-2001
		SE 517136 C2	16-04-2002
		SE 0103046 A	14-11-2001
		US 2001031784 A1	18-10-2001

EPO FORM P0439

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82